

**Early Hearing Detection and Intervention (EHDI)
Region IV Programs
Risk Factor Monitoring Survey**

Primary Reference: Early Hearing Detection and Intervention
Program Guidance Manual
<http://www.cdc.gov/ncbddd/ehdi/documents/EDHI%20Guidelines.pdf>

Appendix III
Surveillance Data Items

CDC Recommended:

M = minimum (recommended for all state data systems; required for follow-up on universal NBHS and for full r
C = core (recommended for complete state data systems
E = enhanced (additional data useful for clinicians, enhanced tracking, or research)

Item	Need
Family History	M
In utero / Congenital infections	M
" CMV	M
" Rubella	M
" Syphilis	M
" Herpes	M
" Toxoplasmosis	M
" Other?	M
Low birth weight	C
" <1500 g	C
APGAR score 0-4 @ 1min or 0-6 @ 5 min	C
Birth Defects	
" Craniofacial (including those with morphologic	M
" Head and neck (e.g., cleft palate, ear tags)	C
" Findings associated with a syndrome known to	M
NICU admission	M
" # days (> 48 h)	M
" < 48 h	C
" 48 h to 28 d	C
Neonatal Indicators	C
" hyperbilirubinemia requiring exchange transfus	M
" Mechanical ventilation (5+ d)	C
" sistent pulmonary hypertension assoc w/mech v	C
" Conditions requiring ECMO	M
Neurodegenerative disorder	C
Ototoxic drugs (specify/list)	C
Bronchio-pulmonary dysplasia	C
Other risk factors at discharge	C
Parent or caregiver concern regarding hearing, speech, language , d	M
Stigmata or other findings associated with SNHL or conductive HL or	M
Postnatal infection	
" Bacterial meningitis	M
" Other associated w/HL	M
Syndromes associate (e.g., Neurofibromatosis (NF) 1 and 2, Usher S	M

Neurodegenerative di (e.g., Hunter sx, Charcot-Marie-Tooth)	M
Head trauma	M
Recurrent or persistent otitis media with effusion (OME) for at least 3	M
Other risk factors ider (describe)	E
Description	
" of syndromes and stigmata	C
" of neurologic disorders	C
" of infections	C
" of parental concerns	C
" of other risk factors	C

Reporting on national EMDI goals)